

Specimen No: MW-35874-1C
Pet/Owner: Brandy / Hernandez
Services: 2T
Serial No: 224624
Expires: 05/15/2010

Veterinarian	Account Number
APRIL GAST	
FOUR SEASONS ANIMAL	3378
HOSPITAL	
10624 RANDOLPH STREET	
CROWN POINT, IN 46307 US	

Set 1 Groups: GBE TBM TCF WDM TEM GJO TOM TPH WPM TSM GTI
Set 2 Groups: MAS SBL MCA MHO SMM SFL MFU MGS MRN

For double treatment sets, Set 1 and Set 2 should be given concurrently, but with separate needles and injection sites.

Vial	Injection Number	Dose (cc)	Days Between Injections	Date of Injection/Comments
<b style="color: green;">GREEN CAP Vial A 1:20,000 wt/vol (min 21 days / max 28 days)	1	0.10	3-4	
	2	0.20	3-4	
	3	0.30	3-4	
	4	0.40	3-4	
	5	0.55	3-4	
	6	0.70	3-4	
	7	0.85	3-4	
<b style="color: blue;">BLUE CAP Vial B 1:2000 wt/vol (min 32 days / max 56 days)	8	0.05	4-7	
	9	0.10	4-7	
	10	0.20	4-7	
	11	0.35	4-7	
	12	0.50	4-7	
	13	0.60	4-7	
	14	0.70	4-7	
	15	0.85	4-7	
<b style="color: red;">RED CAP Vial C 1:200 wt/vol (min 204 days / max 227 days) * Contact your veterinarian to order your treatment refill at this time.	16	0.05	7-8	
	17	0.10	7-8	
	18	0.20	7-8	
	19	0.30	7-8	
	20	0.40	7-8	
	21	0.50	7-8	
	22	0.60	7-8	
	23	0.70	14-21	
	24	0.75	21-30	
	25	0.75	30	
	26	0.75	30	
	27 *	0.75	30	
28	0.75	30		

Before administering the first injection, carefully read the enclosed booklet "Immunotherapy: A Guide to Treating Your Pet's Allergies" and the detailed instructions on the back side of this page.

To increase the comfort level for your pet, a weight-appropriate dose of oral antihistamine should be given 45-60 minutes prior to each allergy injection.

During peak allergy season, more frequent injections may be required, thereby reducing the number of days and/or weeks between injections.