

Specimen No: NE-27884-4E
Pet/Owner: Blue Lagoon / Montgomery
Services: 2TEQ
Serial No: 224606
Expires: 05/15/2010

Veterinarian	Account Number
MICHELE L. FERRARO	
MILLBROOK EQUINE VET	3553
CLINIC	
109 N. MABBETTSVILLE ROAD	
MILLBROOK, NY 12545 US	

Set 1 Groups: TBB TRB GBJ TBM TCX WGL THZ SME TOM GOR GRF SWO
GTI

Set 2 Groups: MAS SBL MGS MHE SHR STP

For double treatment sets, Set 1 and Set 2 should be given concurrently, but with separate needles and injection sites.

Vial	Injection Number	Dose (cc)	Days Between Injections	Date of Injection/Comments
GREEN CAP Vial A 1:1750 wt/vol (min 18 days / max 24 days)	1	0.20	3-4	
	2	0.40	3-4	
	3	0.60	3-4	
	4	0.80	3-4	
	5	1.00	3-4	
	6	1.00	3-4	
BLUE CAP Vial B 1:175 wt/vol (min 24 days / max 42 days)	7	0.20	4-7	
	8	0.40	4-7	
	9	0.60	4-7	
	10	0.80	4-7	
	11	1.00	4-7	
	12	1.00	4-7	
RED CAP Vial C 1:35 wt/vol (min 183 days / max 203 days) * Contact your veterinarian to order your treatment refill at this time.	13	0.10	7-8	
	14	0.20	7-8	
	15	0.40	7-8	
	16	0.60	7-8	
	17	0.80	14-21	
	18	1.00	21-30	
	19	1.00	30	
	20	1.00	30	
	21 *	1.00	30	
	22	1.00	30	

Before administering the first injection, carefully read the enclosed booklet "Immunotherapy: A Guide to Treating Your Pet's Allergies" and the detailed instructions on the back side of this page.

To increase the comfort level for your pet, a weight-appropriate dose of oral antihistamine should be given 45-60 minutes prior to each allergy injection.

During peak allergy season, more frequent injections may be required, thereby reducing the number of days and/or weeks between injections.